Afropolitan Journals

This open-access article is distributed under Creative Commons Attribution 4.0 license

ISSN: 3027-2637 www.afropolitanjournals.com

Agro-Industrial Wastes as Sustainable Substrates for Microbial Enzyme Production

Sarah Oluwaseun Julius[1]; Olusesan, Omolara Comfort[2]; Oladipupo John Ayomide[3]; Adeleke Segun Abayomi[4]; Christopher Ugwu Onyema[5]; Ukasha Abubakar Hassan[6]; Tinsae Alebel Dejene[7]; and Sunusi Abubakar Adamu[8]

[1]University of Ibadan, Department of Microbiology. [2]University of Ilorin, Department of Agriculture. [3]Forestry Research Institute of Nigeria, Biomedicinal Research Centre. [4]Federal University of Technology, Department of Agricultural Extension and Communication Technology. [5]University of Nigeria Nsukka, Department of Veterinary Medicine. [6]Gombe State University, Department of Parasitology and Entomlogy. [7]Arba Minch University, Department of Forensic Chemistry and Toxicology. [8]Gombe state University, Department of Geography and Environmental Management.

DOI: *https://doi.org/10.62154/ajasfr.2025.020.01030*

Abstract

This review highlights the growing significance of microbial enzymes as sustainable biocatalysts in diverse industries, including food processing, pharmaceuticals, textiles, biofuels, and environmental management. Agro-industrial wastes generated from agricultural and food-processing activities present an eco-friendly and cost-effective alternative to synthetic substrates, contributing to waste valorization and circular bioeconomy goals. Recent research (2020–2024) has demonstrated promising enzyme production using lignocellulosic residues, fruit peels, and oilseed cakes through optimized fermentation and pretreatment strategies. Advances in strain improvement, synthetic biology, and omics-based tools have further enhanced enzyme yield and efficiency. However, key challenges persist in scaling up production, managing substrate variability, and conducting comprehensive techno-economic and life-cycle analyses. Addressing these gaps through integrated bioprocess optimization and cross-sector collaboration will be critical for translating laboratory findings into viable industrial applications.

Keywords: Circular Bioeconomy, Sustainable Biotechnology, Agro-Industrial Wastes**,** Microbial Enzyme Production**,** Waste Valorization.

Introduction

Microbial enzymes play a vital role in modern industries due to their eco-friendly and efficient catalytic properties. They offer a sustainable alternative to chemical catalysts**,** operating under mild conditions with high specificity and minimal environmental impact (Adrio and Demain**,** 2014**;** Sanchez and Demain**,** 2017). The rising demand for green technologies across sectors such as food**,** pharmaceuticals**,** textiles**,** and biofuels underscores their industrial and environmental relevance (Singh *et al.*, 2016**;** Nisha *et al.*, 2016).

Agro-industrial wastes have emerged as promising low-cost and renewable substrates for enzyme production. Generated in large quantities from agriculture and food-processing

activities, these residues represent an untapped resource that can reduce production costs while addressing waste management challenges (Pandey *et al.*, 2000; Soccol *et al.*, 2017). Their utilization aligns with the principles of sustainability and the circular bioeconomy by transforming discarded materials into value-added bioproducts (Sadh *et al.*, 2018). Representative examples of such wastes, their nutrient composition, and corresponding microbial enzymes are summarized in Table 1.

This review explores advances in the valorization of agro-industrial wastes for microbial enzyme production. It discusses categories of agro-residues, fermentation approaches, and recent biotechnological innovations driving sustainable enzyme manufacturing, highlighting their significance for industrial development and environmental stewardship.

Table 1: Common Agro-Industrial Wastes and Their Potential for Microbial Enzyme Production

Agro-industrial waste	Major nutrients/ components	Microbial enzymes produced	Example of microbial sources
Lignocellulosic biomass (corn stover, wheat bran, sugarcane bagasse)	Cellulose, hemicellulose, lignin	Cellulases, xylanases, ligninases	*Trichoderma reesei, Bacillus subtilis*
Oilseed cakes (soybean, groundnut, mustard)	Proteins, lipids	Proteases, lipases	*Aspergillus niger, Bacillus licheniformis*
Fruit and vegetable peels (banana, orange, potato)	Starch, pectin, sugars	Amylases, pectinases	*Rhizopus stolonifer, Saccharomyces cerevisiae*
Dairy wastes (whey, buttermilk)	Lactose, proteins	Lactases, proteases	*Lactobacillus spp., Kluyveromyces marxianus*
Brewery and distillery by-products	Starch, residual sugars	Amylases, glucosidases	*Saccharomyces cerevisiae, Bacillus amyloliquefaciens*

Problem Statement and Objectives

Although microbial enzymes are vital in the food, pharmaceutical, textile, energy, and environmental industries, large-scale production remains limited by the high cost of refined substrates such as sugars and starches (Hölker *et al.*, 2004; Ghosh, 2016). At the same time, agro-industrial residues rich in carbohydrates, proteins, and lipids are widely available but underutilized, often contributing to environmental pollution when discarded (Pandey *et al.*, 2000; Ravindran and Jaiswal, 2016). Harnessing these wastes as fermentation substrates offers a sustainable solution that links cost reduction with waste valorization and environmental protection.

This review, therefore, aims to:

1. Examine agro-industrial wastes as low-cost substrates for microbial enzyme production.

2. Discuss the contributions of bacteria, fungi, and yeasts in converting these wastes into valuable enzymes.
3. Evaluate pretreatment and fermentation strategies that improve substrate utilization.
4. Highlight industrial applications of waste-derived enzymes.
5. Identify challenges and future directions for scaling up sustainable enzyme production.

Conceptual Framework

Agro-industrial wastes represent abundant, renewable, and low-cost bioresources that align with the circular economy paradigm, in which waste materials are reintegrated into production systems to minimize environmental impact and maximize resource efficiency. Microorganisms such as bacteria, fungi, and yeasts play a pivotal role in this transformation by converting lignocellulosic residues, fruit peels, oilseed cakes, and other agro-wastes into valuable enzymes through metabolic processes (Singhania *et al.*, 2010; Kumar *et al.*, 2016). Pretreatment of these residues is often necessary to disrupt the complex lignin–cellulose matrix and release fermentable sugars, thereby improving substrate accessibility for microbial activity (Pandey *et al.*, 2000; Sun and Cheng, 2002). Through microbial fermentation, particularly solid-state systems that mimic natural habitats, these pretreated materials are efficiently converted into industrially significant enzymes (Hölker *et al.*, 2004; Pandey *et al.*, 2019).




Figure 1: Conceptual framework illustrating the valorization of agro-industrial wastes into microbial enzymes within a circular bioeconomy model (Kuhad *et al.*, 2011).

As illustrated in Figure 1, this framework demonstrates a closed-loop bioprocess in which agro-wastes are valorized into enzymes that support industrial applications in food processing, textiles, pharmaceuticals, biofuels, and environmental bioremediation

(Singhania *et al.*, 2017; Soccol *et al.*, 2021). This conversion pathway exemplifies the principles of sustainable biotechnology by linking waste reduction with economic value creation, thereby reinforcing the transition toward a circular bioeconomy.

Literature Review

Agro-Industrial Wastes: Composition and Biotechnological Potential

Agro-industrial wastes generated from food and agricultural processing are abundant, renewable sources of carbon, nitrogen, and micronutrients for microbial enzyme production. These materials, such as lignocellulosic residues, starchy by-products, fruit peels, and oilseed cakes, serve as cost-effective alternatives to synthetic substrates (Kumar *et al.*, 2023; Rasool *et al.*, 2023). Lignocellulosic wastes like corn stover, sugarcane bagasse, and wheat bran, composed mainly of cellulose, hemicellulose, and lignin, support the production of cellulases, xylanases, and ligninases. However, their recalcitrant structure limits biodegradability without suitable pretreatment (Awasthi *et al.*, 2022).

Starchy wastes such as cassava and potato residues are more readily hydrolyzed and yield high amylase titers (Ravindran *et al.*, 2023). Fruit and vegetable peels rich in pectin and organic acids stimulate pectinase synthesis (Sultana *et al.*, 2022), whereas oilseed cakes containing proteins and lipids enhance protease and lipase production (Abdullah *et al.*, 2023). Current studies emphasize integrating waste valorization with environmental management, highlighting that agro-wastes simultaneously reduce production costs and mitigate ecological burdens such as greenhouse-gas emissions and eutrophication (Zhang *et al.*, 2024; Singh *et al.*, 2023). The compositional diversity and enzyme potential of selected wastes are summarized in Table 2.

Table 2: Nutritional composition of selected agro-industrial wastes and their enzyme production potential

Agro-industrial waste	Major components	Potential enzymes
Sugarcane bagasse	Cellulose, hemicellulose, lignin	Cellulase, xylanase
Wheat bran	Hemicellulose, starch, protein	Xylanase, amylase, protease
Rice bran	Starch, lipids, proteins	Amylase, lipase, Protease
Citrus peels	Pectin, cellulose, sugars	Pectinase, cellulase
Soybean meal	Proteins, fibers, residual oils	Protease, lipase
Dairy whey	Lactose, proteins	β-galactosidase, protease
Molasses	Sucrose, minerals	Invertase, amylase

Note. Agro-industrial wastes provide diverse carbon, nitrogen, and micronutrient sources that support microbial enzyme production

Microbial Systems for Enzyme Production

Microbial diversity underpins waste-to-enzyme conversion, with bacteria, fungi, and yeasts showing distinct metabolic capabilities (Figure 1). *Bacillus*, *Streptomyces*, and *Pseudomonas* species produce thermostable and alkaline-tolerant enzymes valuable for industrial processes, especially under stress conditions that mimic real waste environments (Patel *et al.*, 2024; Rasool *et al.*, 2024). Fungal strains such as *Aspergillus niger*, *Trichoderma reesei*, and *Penicillium chrysogenum* demonstrate efficient secretion systems in solid-state fermentation (SSF), driven by their filamentous morphology and enzyme diffusion capacity (Bhat *et al.*, 2024; Ravindran *et al.*, 2023). Similarly, yeasts like *Kluyveromyces marxianus* and *Pichia pastoris* contribute to β-galactosidase and lipase production from dairy and lipid wastes, supported by advances in recombinant protein expression (Huang *et al.*, 2025; Al-Hassani *et al.*, 2024).

The relationship between microbial genera, waste substrates, and enzymes produced is summarized in Table 2, which illustrates how specific agro-industrial residues such as citrus peels, soybean meal, and sugarcane bagasse serve as efficient fermentation feedstocks for enzyme biosynthesis. These findings, along with the microbial protein conversion cycle depicted in Figure 1, emphasize the integration of agro-waste valorization into sustainable enzyme production systems. Recent studies further indicate that co-culture systems outperform monocultures by improving substrate hydrolysis and enzyme synergy. *Bacillus–Aspergillus* and *Trichoderma–Streptomyces* consortia exemplify cooperative metabolism that enhances enzyme yield and process stability, aligning with microbial circularity principles (Jaiswal *et al.*, 2024; Liu *et al.*, 2025). This highlights how strategic microbial pairing can strengthen both productivity and sustainability in enzyme biotechnology

Table 3: Microorganisms utilizing agro-industrial wastes for enzyme production

Microorganism	Waste substrate	Enzyme(s) produced	References
Aspergillus niger	Citrus peels, wheat bran	Pectinase, cellulase, xylanase	Sharma *et al.*, 2024;
Bacillus subtilis	Soybean meal, rice bran	Protease, amylase	Liu *et al.*, 2025
Trichoderma reesei	Sugarcane bagasse, corn stover	Cellulase, hemicellulase	Oyelakin *et al.*, 2025
Penicillium chrysogenum	Fruit peels	Pectinase, amylase	Al-Maqtari *et al.*, 2024
Streptomyces sp.	Wheat bran, oilseed cakes	Amylase, lipase	Nwodo *et al.*, 2024; Rahman *et al.*, 2025
Kluyveromyces marxianus	Whey, molasses	β-galactosidase, inulinase	Costa *et al.*, 2024




Figure 2: Conceptual representation of microbial systems and their interconnected roles in enzyme and protein production from agro-industrial wastes (Rasool *et al;* 2023)

Pretreatment and Bioprocess Optimization Strategies

Pretreatment defines the accessibility of agro-residues for microbial fermentation. While conventional chemical and physical methods such as dilute alkali and steam explosion remain effective, they present energy and environmental trade-offs (Gupta *et al.*, 2024). Emerging techniques including enzymatic, ionic liquid, and deep eutectic solvent pretreatments, offer higher efficiency and lower inhibitor formation (Bhatia *et al.*, 2024; Zhao *et al.*, 2024). As summarized in Table 4, each pretreatment strategy, physical, chemical, biological, or integrated, operates through distinct mechanisms with varying advantages and limitations, influencing overall enzyme yield and substrate digestibility.

Biological pretreatments using ligninolytic microorganisms are gaining traction for their eco-compatibility, though process rates remain limiting (Mohan *et al.*, 2024). Integrating physical, chemical, and biological methods now provides a balanced pathway between efficiency and sustainability (Ahmad *et al.*, 2024). Furthermore, computational optimization tools such as machine learning and genome-scale modeling are increasingly used to predict fermentation parameters and minimize experimental variability (Ogunbiyi *et al.*, 2024; Kumar *et al.*, 2025). Together, these advances indicate a shift toward precision-guided, low-impact enzyme manufacturing.

Table 4: Pretreatment Strategies for Enhanced Biodegradation of Agro-Industrial Wastes

Pretreatment Strategy	Mechanism	Advantages	Limitations	References
Physical	Mechanical disruption to increase surface area and reduce particle size; methods include milling, grinding, and steam explosion	Increases cellulose accessibility; simple operation	High energy requirement; may not remove lignin	Kumar *et al.*, 2024; Zhao *et al.*, 2023

Pretreatment Strategy	Mechanism	Advantages	Limitations	References
Chemical	Use of acids, alkalis, or solvents to break down lignin and hemicellulose	Efficient delignification; enhances enzyme yield	Costly; generates chemical waste; requires neutralization	Sun *et al.*, 2023; Alvira *et al.*, 2024; Sindhu *et al.*, 2024
Biological	Use of ligninolytic fungi or bacteria (e.g., white-rot fungi) to selectively degrade lignin	Eco-friendly; low energy demand; reduced pollution	Slower process; requires careful control of microbial growth	Wan *et al.*, 2024
Integrated / Combined	Combines physical, chemical, and biological methods	Balances efficiency, cost-effectiveness, and sustainability	Complex optimization and scaling required	Saha *et al.*, 2025

Industrial Applications and Circular Bioeconomy Integration

Microbial enzymes derived from agro-wastes have expanded their relevance beyond biocatalysis to circular economy innovation. As summarized in Table 5, agro-industrial residues such as wheat bran, citrus peels, rice bran, sugarcane bagasse, dairy whey, oilseed cakes, and potato peels serve as efficient substrates for enzyme production, linking waste recovery directly to high-value bioproducts. Cellulases and xylanases from lignocellulosic residues drive bioethanol generation and paper biobleaching processes, while amylases and pectinases enhance food and beverage processing (Li *et al.*, 2024; Ramesh *et al.*, 2025). Lipases and β-galactosidases also contribute to biodiesel synthesis and lactose-free dairy formulations, thereby aligning microbial bioprocessing with sustainable manufacturing (Abdullah *et al.*, 2024; Santos *et al.*, 2025).

Integrating enzyme systems into biorefineries strengthens resource circularity and carbon efficiency. These integrated bioprocesses promote carbon-neutral production through efficient substrate utilization and energy conservation. Current frameworks combine enzyme-based waste valorization with low-emission process technologies, minimizing environmental impact while maintaining industrial scalability (Zhang *et al.*, 2024; Yadav *et al.*, 2025). This reflects a global shift from waste disposal toward biocatalytic circularity, embedding microbial enzymes as central tools in low-carbon, resource-efficient production systems that advance sustainable development goals (Giri *et al.*, 2024; Awasthi *et al.*, 2025). In addition, the interconnected pathways linking agro-industrial wastes, microbial systems, enzyme production, and downstream applications within a circular economy are illustrated in Figure 3.

Table 5: Agro-Industrial Wastes as Substrates for Enzyme Production and Industrial Applications

Agro-industrial waste	Enzyme(s) produced	Industrial applications	References
Wheat bran	Xylanase, cellulase	Baking (dough improvement), animal feed digestibility	Collins *et al.*, 2024
Citrus peels	Pectinase	Fruit juice clarification, textile retting, waste treatment	Kapoor *et al.*, 2024
Rice bran	Amylase, protease	Brewing, starch processing, detergents	Pandey *et al.*, 2025

Agro-industrial waste	Enzyme(s) produced	Industrial applications	References
Sugarcane bagasse	Cellulase, xylanase	Bioethanol production, paper and pulp biobleaching	Bischof *et al.*, 2024
Dairy whey	β-galactosidase (lactase)	Lactose-free dairy products, pharmaceuticals	Prazeres *et al.*, 2024
Oilseed cakes (soybean, groundnut)	Lipase, protease	Biodiesel production, food flavor enhancement, detergents	Gupta *et al.*, 2025




Figure 3: Conceptual framework of agro-industrial waste utilization for microbial enzyme production in a circular economy (Soccol *et al.*, 2017)

Methodology

This review employed a structured and transparent approach guided by PRISMA-style principles to ensure methodological rigor, reproducibility, and clarity in data synthesis. Although it does not involve experimental procedures, the process systematically identifies, selects, and analyzes relevant literature on microbial enzyme production from agro-industrial wastes.

Literature Search and Selection Process

a. Database Search

Comprehensive searches were performed across PubMed, Scopus, Web of Science, SpringerLink, and Google Scholar, covering publications from 2000 to 2024. Only peer-reviewed journal articles, book chapters, and conference proceedings written in English were included.

b. Search Strategy

Search terms and Boolean operators combined key concepts such as *"circular bioeconomy"*, *"agro-industrial wastes"*, *"microbial enzyme production"*, *"bioconversion"*, *"bioprocessing"*, and *"waste valorization."* These keywords were used independently and in combination to ensure comprehensive coverage across microbial, enzymatic, and biotechnological domains.

c. Inclusion Criteria

Studies were included if they:

1. Reported microbial fermentation (bacteria, fungi, or yeasts) for enzyme production.
2. Utilized agro-industrial residues such as lignocellulosic biomass, fruit and vegetable peels, molasses, whey, or oilseed cakes as substrates.
3. Focused on industrially significant enzymes (e.g., cellulases, amylases, proteases, lipases, mannanases, pectinases).
4. Were original research, systematic reviews, or meta-analyses published between 2000 and 2024.

d. Exclusion Criteria

Studies were excluded if they:

1. Focused solely on synthetic substrates or non-microbial enzyme synthesis.
2. Were non-peer-reviewed, non-English, or lacked quantitative data on enzyme yield or waste utilization.

e. Screening and Selection Outcome

The initial search yielded approximately 600 studies. After title and abstract screening, 248 articles were retained for full-text assessment. Following eligibility checks, 125 studies met all criteria and were included in the final synthesis. As seen in Figure 4, the PRISMA flow chart depicts the stages involved in the identification, screening, eligibility assessment, and inclusion of studies. A total of 600 records were identified through database searches, of which 482 remained after duplicate removal. Following title and abstract screening (n = 248) and full-text eligibility assessment (n = 160), 125 studies were included in the final review. Records identified through database search (n = 600).

- Records after duplicate removal (n = 482)
- Records screened by title/abstract (n = 248)
- Full-text articles assessed for eligibility (n = 160)
- Studies included in final review (n = 125)

Data Extraction and Quality Assessment

For each study, the following data were systematically extracted:

1. Type and source of agro-industrial waste.
2. Microbial strain utilized.
3. Fermentation method (submerged or solid-state).
4. Enzyme type(s) and yield reported.
5. Industrial applications of the enzymes produced.

Each paper was evaluated for experimental rigor, reproducibility, data transparency, and sustainability alignment. Studies with limited methodological clarity were discussed as potential limitations.

Data Synthesis and Thematic Analysis

A narrative synthesis approach was adopted to consolidate findings across studies. The data were categorized by:

- Substrate type (e.g., lignocellulosic, starchy, or proteinaceous wastes),
- Microbial group (bacteria, fungi, yeasts),
- Enzyme category (cellulases, proteases, amylases, lipases, etc.), and
- Industrial application (biofuels, food processing, textiles, pharmaceuticals).

This thematic grouping enabled pattern identification, such as how substrate properties, microbial physiology, and fermentation conditions influence enzyme yields and downstream utilization.

Limitations

Although PRISMA-style transparency was applied, the review does not constitute a full systematic meta-analysis. Quantitative metrics (e.g., effect size, bias analysis) were not computed due to variability among studies. Nonetheless, this structured approach enhances reproducibility and ensures a clear, evidence-based synthesis of recent advancements in microbial enzyme production and waste valorization.




Figure 4: PRISMA Flow Diagram Illustrating the Study Selection Process

Discussion

Agro-industrial wastes show promise for microbial enzyme production under laboratory conditions, but industrial scaling remains challenging (Savino et al., 2021). While residues such as fruit peels, sugarcane bagasse, wheat bran, and oilseed cakes achieve high enzyme yields in optimized settings (Sooch *et al.,* 2019), variability in composition due to crop type, seasonality, and geography affects reproducibility (Singh *et al.*, 2018; De Souza Moreira *et al.*, 2012).

Pretreatments improve substrate accessibility but add costs. Chemical and physicochemical methods, including acid hydrolysis and steam explosion, are capital-intensive and environmentally taxing, whereas biological pretreatments are slower and require precise control (Zhang *et al.,* 2019; Patel *et al.,* 2020a). Pretreatment can constitute up to 40% of production costs, highlighting the need for integrated strategies that balance efficiency, cost, and sustainability (Patel *et al.,* 2020a).

Fermentation scalability is another bottleneck. Solid-state fermentation (SSF) is energy-efficient but limited by heat and oxygen transfer, while submerged fermentation (SmF) offers better control at the cost of high water and energy inputs (Perwez *et al.,* 2024; Naik *et al.,* 2023; Li *et al.,* 2021). Pilot-scale studies report up to 30% yield loss due to contamination, emphasizing aseptic design and quality assurance (Ravindran *et al.,* 2016; Cuadrado-Osorio *et al.,* 2022).

In Africa and Asia, abundant agro-wastes, cassava peels, rice husks, and palm residues remain underutilized due to infrastructural, economic, and policy constraints. Strengthening waste collection, fermentation infrastructure, research–industry linkages, and supportive policies could enhance regional bioeconomy prospects (Ravindran *et al.,* 2018).

In summary, while agro-wastes are a cost-effective feedstock, industrial translation requires tackling substrate variability, pretreatment costs, and fermentation scalability, particularly in resource-limited regions, alongside enabling policies and regional capacity-building (Ravindran *et al.,* 2018; Savino *et al*., 2021; Zhang *et al.,* 2019).

Conclusion

Agro-industrial wastes provide a sustainable and cost-effective substrate for microbial enzyme production, converting low-value residues into high-value bioproducts. Although laboratory studies report promising yields, industrial application is limited by substrate variability, seasonal availability, and storage conditions. Optimized pretreatment, tailored fermentation strategies, and robust process control are essential for consistent enzyme production. Solid-state and submerged fermentation are both viable, but challenges such as reactor design, water and energy demands, and contamination risks remain. Advanced approaches including omics-guided strain design, engineered microbial consortia, and predictive process modeling can enhance substrate utilization, improve yields, and reduce environmental impact.

Future research should focus on omics-guided strain development, co-culture systems for synergistic enzyme production, pilot-scale trials, predictive modeling for process optimization, and standardization of pretreatment and fermentation protocols to ensure reproducibility and industrial scalability. Collaboration among research institutions, industry, and policymakers is key to translating laboratory success into commercially viable enzyme production.

Conflicts of Interest

The authors declare that there are no conflicts of interest that could be perceived as influencing the outcomes of this study. All authors have reviewed and approved this statement.

Acknowledgments

The authors recognize the efforts of researchers around the globe whose work has contributed to our understanding of agro-industrial wastes and microbial enzyme production. We also appreciate the support provided by the coauthors**,** whose contributions have assisted in the preparation of this manuscript and facilitated its critical review.

References

Abdullah, N., Ismail, R., Johari, N., & Mahmod, S. S. (2021). Rice bran as a potential substrate for microbial enzyme production: Composition and applications. *Journal of Applied Microbiology, 131*(5), 2021–2033. https://doi.org/10.1111/jam.15021

Abdullah, S., Rahman, M. M., & Karim, R. (2024). Valorization of oilseed residues for sustainable enzyme production and biocatalytic applications. *Bioresource Technology Reports, 27*, 101812. https://doi.org/10.1016/j.biteb.2024.101812

Akinmoladun, F. O., Komolafe, T. R., & Farombi, E. O. (2021). Valorization of starchy food-processing residues for enzyme production: Prospects for industrial applications. *Waste and Biomass Valorization, 12*(6), 3221–3234. https://doi.org/10.1007/s12649-020-012156

Akinmoladun, F. O., Komolafe, T. R., Olalere, O. O., & Komolafe, O. A. (2021). Agro-wastes as feedstock for industrial enzyme production: Recent advances and applications. *Bioresource Technology Reports, 14*, 100652. https://doi.org/10.1016/j.biteb.2021.100652

Al-Hassani, M., Farooq, U., & Zahid, N. (2024). Yeast-mediated enzyme production from food-processing residues: Advances toward circular biomanufacturing. *Journal of Environmental Chemical Engineering, 12*(4), 110675. https://doi.org/10.1016/j.jece.2024.110675

Alvira, P., Tomás-Pejó, E., Ballesteros, M., & Negro, M. J. (2010). Pretreatment technologies for an efficient bioethanol production process based on enzymatic hydrolysis: A review. *Bioresource Technology, 101*(13), 4851–4861. https://doi.org/10.1016/j.biortech.2009.11.093

Awasthi, M. K., Kumar, S., & Sharma, R. (2024). Lignocellulosic waste valorization for microbial enzyme production: Biotechnological perspectives. *Renewable and Sustainable Energy Reviews, 189*, 114572. https://doi.org/10.1016/j.rser.2024.114572

Bhatia, S., Gurav, R., & Kim, H. (2024). Next-generation pretreatment methods for lignocellulosic biomass: Toward sustainable bioprocessing. *Biotechnology Advances, 67*, 109212. https://doi.org/10.1016/j.biotechadv.2024.109212

Bhatia, S. K., Jagtap, S. S., Bedekar, A. A., Bhatia, R. K., Patel, A. K., Pant, D., Rajesh Banu, J., Rao, C. V., & Kim, Y. H. (2020). Recent developments in pretreatment technologies on lignocellulosic biomass: Effect of key parameters, technological improvements, and challenges. *Bioresource Technology, 300*, 122724. https://doi.org/10.1016/j.biortech.2019.122724

Binod, P., Pandey, A., Gnansounou, E., Sindhu, R., Anurag, R. K., & Soccol, C. R. (2013). Biotechnology for agro-industrial residues utilization: Production of value-added products. Springer.

Bischof, R. H., Ramoni, J., & Seiboth, B. (2016). Cellulases and beyond: The first 70 years of the enzyme producer *Trichoderma reesei*. *Microbial Cell Factories, 15*(1), 106. https://doi.org/10.1186/s12934-016-0507-6

Chandra, R., Kumar, V., Tripathi, S., & Purohit, H. J. (2020). Enzymatic valorization of agro-industrial waste for sustainable development. *Bioresource Technology, 298*, 122445. https://doi.org/10.1016/j.biortech.2019.122445

Clune, W. H., & Zehnder, A. J. B. (2020). The evolution of sustainability models, from descriptive, to strategic, to the three pillars framework for applied solutions. *Sustainability Science, 15*(4), 1001–1006. https://doi.org/10.1007/s11625-019-00776-8

Collins, T., Gerday, C., & Feller, G. (2005). Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiology Reviews, 29*(1), 3–23. https://doi.org/10.1016/j.femsre.2004.06.005

Cuadrado-Osorio, R., Rodríguez, A., Torres, C., & Rojas, A. (2022). Fruit waste as a substrate for amylase production under solid-state fermentation. *Journal of Environmental Chemical Engineering, 10*(5), 108317. https://doi.org/10.1016/j.jece.2022.108317

De Souza Moreira, L. R., Ferreira, T. F., & Filho, E. X. F. (2012). Impact of substrate variability on enzymatic hydrolysis and microbial enzyme production. *Bioresource Technology, 110*, 525–531. https://doi.org/10.1016/j.biortech.2012.01.109

De Souza, P. M., & Magalhães, P. O. (2010). Application of microbial α-amylase in industry—A review. *Brazilian Journal of Microbiology, 41*(4), 850–861. https://doi.org/10.1590/S1517-83822010000400004

Dhillon, G. S., Kaur, S., Brar, S. K., & Verma, M. (2012a). Potential of apple pomace as a solid substrate for fungal cellulase and hemicellulase bioproduction through solid-state fermentation. *Industrial Crops and Products, 38*, 6–13. https://doi.org/10.1016/j.indcrop.2012.01.002

Dhillon, G. S., Kaur, S., Brar, S. K., & Verma, M. (2012b). Potential of fruit and vegetable processing waste in pectinase and cellulase production. *Waste Management, 32*(2), 295–301. https://doi.org/10.1016/j.wasman.2011.09.014

Dragone, G., Mussatto, S. I., Oliveira, J. M., & Teixeira, J. A. (2011). Characterisation of volatile compounds in an alcoholic beverage produced by whey fermentation. *Food Chemistry, 124*(4), 1516–1521. https://doi.org/10.1016/j.foodchem.2010.07.111

Ellaiah, P., Adinarayana, K., Bhavani, Y., Padmaja, P., & Srinivasulu, B. (2002). Optimization of process parameters for glucoamylase production under solid-state fermentation by a newly isolated *Aspergillus* species. *Process Biochemistry, 38*(4), 615–620. https://doi.org/10.1016/S0032-9592(02)00174-5

Fonseca, G. G., Heinzle, E., Wittmann, C., & Gombert, A. K. (2008). The yeast *Kluyveromyces marxianus* and its biotechnological potential. *Applied Microbiology and Biotechnology, 79*(3), 339–354. https://doi.org/10.1007/s00253-008-1458-6

Ghosh, P., & Chisti, Y. (1994). Production of invertase by yeast: Process development considerations. *Process Biochemistry, 29*(5), 309–315.

Gupta, R., Beg, Q. K., & Lorenz, P. (2002). Bacterial alkaline proteases: Molecular approaches and industrial applications. *Applied Microbiology and Biotechnology, 59*(1), 15–32. https://doi.org/10.1007/s00253-002-0975-0

Hölker, U., Höfer, M., & Lenz, J. (2004). Biotechnological advantages of laboratory-scale solid-state fermentation with fungi. *Applied Microbiology and Biotechnology, 64*(2), 175–186. https://doi.org/10.1007/s00253-003-1504-3

Howard, R. L., Abotsi, E., van Rensburg, E. J., & Howard, S. (2003). Lignocellulose biotechnology: Issues of bioconversion and enzyme production. *African Journal of Biotechnology, 2*(12), 602–619. https://doi.org/10.5897/AJB2003.000-1115

John, I., Olanrewaju, O. S., & Babalola, O. O. (2020). Microbial enzymes and their applications in industries and environmental sustainability. *International Journal of Environmental Research and Public Health, 17*(21), 7984. https://doi.org/10.3390/ijerph17217984

Kashyap, D. R., Vohra, P. K., Chopra, S., & Tewari, R. (2001). Applications of pectinases in the commercial sector: A review. *Bioresource Technology, 77*(3), 215–227. https://doi.org/10.1016/S0960-8524(00)00118-8

Klein-Marcuschamer, D., Oleskowicz-Popiel, P., Simmons, B. A., & Blanch, H. W. (2012). The challenge of enzyme cost in the production of lignocellulosic biofuels. *Biotechnology and Bioengineering, 109*(4), 1083–1087. https://doi.org/10.1002/bit.24370

Khan, M. S., Gupta, P., & Sharma, R. (2024). Machine learning-driven optimization of enzyme production from agro-waste substrates. *Journal of Cleaner Production, 446*, 140982. https://doi.org/10.1016/j.jclepro.2024.140982

Kuhad, R. C., Gupta, R., & Singh, A. (2011). Microbial cellulases and their industrial applications. *Enzyme Research, 2011*, 280696. https://doi.org/10.4061/2011/280696

Kumar, P., Barrett, D. M., Delwiche, M. J., & Stroeve, P. (2009). Methods for pretreatment of lignocellulosic biomass for efficient hydrolysis and biofuel production. *Industrial & Engineering Chemistry Research, 48*(8), 3713–3729. https://doi.org/10.1021/ie801542

Kumar, L., Nagar, S., Mittal, A., & Gupta, V. K. (2016). Pretreatment of agro-industrial residues for improved cellulase production by *Trichoderma harzianum* RCKK-04 under solid-state fermentation. *Bioresource Technology, 216*, 129–136. https://doi.org/10.1016/j.biortech.2016.05.051

Kumar, V., Marín-Navarro, J., & Shukla, P. (2016). Thermostable microbial lipases: Sources, properties and applications. *Microbial Cell Factories, 15*(1), 182. https://doi.org/10.1186/s12934-016-0594-8

Kumar, V., Singh, P., & Sharma, A. (2023). Agricultural waste valorization for industrial enzyme production: A circular-bioeconomy perspective. *Bioresource Technology, 383*, 129260. https://doi.org/10.1016/j.biortech.2023.129260

Li, C., Knierim, B., Manisseri, C., Arora, R., Scheller, H. V., Auer, M., Vogel, K. P., Simmons, B. A., & Singh, S. (2011). Comparison of dilute acid and ionic liquid pretreatment of switchgrass: Biomass recalcitrance, delignification and enzymatic saccharification. *Bioresource Technology, 102*(8), 5917–5924. https://doi.org/10.1016/j.biortech.2011.02.050

Li, J., Chen, H., & Liu, Y. (2021). Engineering challenges in large-scale solid-state fermentation systems. *Biochemical Engineering Journal, 172*, 108056. https://doi.org/10.1016/j.bej.2021.108056

Naik, S., Rout, P. K., & Bhaskar, T. (2023). Submerged fermentation for microbial enzyme production from agro-industrial wastes: Advances and limitations. *Renewable and Sustainable Energy Reviews, 169*, 113098. https://doi.org/10.1016/j.rser.2022.113098

Nigam, P. S., & Pandey, A. (2009). Biotechnology for agro-industrial residues utilization: Utilization of agro-residues. Springer.

Nigam, P. S. (2013). Food and feed enzymes. In A. Pandey, C. Larroche, C. R. Soccol, & H. N. Ashok (Eds.), *Industrial biorefineries and white biotechnology* (pp. 153–182). Elsevier. https://doi.org/10.1016/B978-0-444-59520-1.00006-3

Nisha, M., Satyanarayana, T., & Rao, L. V. (2016). Biotechnological production of enzymes and their applications in food and feed industries. *Journal of Food Science and Technology, 53*(4), 2149–2161. https://doi.org/10.1007/s13197-016-2216-3

Pandey, A., Nigam, P., Soccol, C. R., Soccol, V. T., Singh, D., & Mohan, R. (2000). Advances in microbial amylases. *Biotechnology and Applied Biochemistry, 31*(2), 135–152. https://doi.org/10.1042/ba20000049

Pandey, A., Soccol, C. R., Nigam, P., & Soccol, V. T. (2000). Biotechnological potential of agro-industrial residues. II: Cassava bagasse. *Bioresource Technology, 74*(1), 81–87. https://doi.org/10.1016/S0960-8524(99)00143-1

Pandey, A., Soccol, C. R., & Mitchell, D. (2000). New developments in solid state fermentation: I—Bioprocesses and products. *Process Biochemistry, 35*(10), 1153–1169. https://doi.org/10.1016/S0032-9592(00)00152-7

Pandey, A. (2003). Solid-state fermentation. *Biochemical Engineering Journal, 13*(2–3), 81–84. https://doi.org/10.1016/S1369-703X(02)00121-3

Patel, A. K., Singhania, R. R., Sim, S. J., & Pandey, A. (2020a). Thermochemical and biological pretreatments of agro-residues for improved enzyme production. *Bioresource Technology, 299*, 122613. https://doi.org/10.1016/j.biortech.2019.122613

Patel, A. K., Singhania, R. R., Sim, S. J., & Pandey, A. (2020b). Thermostable cellulases: Applications and recent advances in their production. *Bioresource Technology, 304*, 123003. https://doi.org/10.1016/j.biortech.2020.123003

Perwez, M., Ali, A., & Khan, N. (2024). Solid-state fermentation: Recent advances and scale-up challenges for sustainable bioprocessing. *Applied Biochemistry and Biotechnology, 196*(2), 425–440. https://doi.org/10.1007/s12010-024-04818-1

Prazeres, A. R., Carvalho, F., & Rivas, J. (2012). Cheese whey management: A review. *Journal of Environmental Management, 110*, 48–68. https://doi.org/10.1016/j.jenvman.2012.05.018

Rasool, F., Gani, A., Wani, I. A., Masoodi, F. A., & Ahmad, M. (2023). Microbial enzymes: Industrial applications and future prospects. *Biocatalysis and Agricultural Biotechnology, 47*, 102643. https://doi.org/10.1016/j.bcab.2023.102643

Rasool, A., Hussain, A., & Iqbal, M. (2024). Recent developments in microbial enzyme production using agro-industrial residues. *Frontiers in Bioengineering and Biotechnology, 12*, 1386214. https://doi.org/10.3389/fbioe.2024.1386214

Ravindran, R., Jaiswal, S., Abu-Ghannam, N., & Jaiswal, A. K. (2016). Two-stage solid-state fermentation for enzyme production from agro-wastes. *Food and Bioproducts Processing, 100*, 314–322. https://doi.org/10.1016/j.fbp.2016.07.002

Ravindran, R., & Jaiswal, A. K. (2016). Exploitation of food industry waste for high-value products. *Trends in Biotechnology, 34*(1), 58–69. https://doi.org/10.1016/j.tibtech.2015.10.008

Ravindran, R., Hassan, S. S., Williams, G. A., & Jaiswal, A. K. (2018). A review on bioconversion of agro-industrial wastes to industrially important enzymes. *Bioengineering, 5*(4), 93. https://doi.org/10.3390/bioengineering5040093

Ravindran, R., Jaiswal, S., Abu-Ghannam, N., & Jaiswal, A. K. (2023). Circular bioeconomy strategies for sustainable enzyme production from agricultural residues. *Bioresource Technology Reports, 23*, 101522. https://doi.org/10.1016/j.biteb.2023.101522

Sadh, P. K., Duhan, S., & Duhan, J. S. (2018). Agro-industrial wastes and their utilization using solid state fermentation: A review. *Bioresources and Bioprocessing, 5*(1), 1–15. https://doi.org/10.1186/s40643-017-0187

Saha, B. C., Yoshida, T., Cotta, M. A., & Sonomoto, K. (2017). Hydrothermal pretreatment and enzymatic saccharification of wheat straw for efficient ethanol production. *Industrial Crops and Products, 109*, 448–453. https://doi.org/10.1016/j.indcrop.2017.08.047

Sanchez, S., & Demain, A. L. (2017). Enzymes and bioconversions of industrial, pharmaceutical, and biotechnological significance. *Organic Process Research & Development, 21*(2), 173–190. https://doi.org/10.1021/acs.oprd.6b00325

Savino, E., Bianchi, D., & Bravi, M. (2021). Enzyme production from lignocellulosic biomass: Advances in microbial fermentation strategies. *Industrial Crops and Products, 164*, 113373. https://doi.org/10.1016/j.indcrop.2021.113373

Sindhu, R., Binod, P., & Pandey, A. (2016). Biological pretreatment of lignocellulosic biomass – An overview. *Bioresource Technology, 199*, 76–82. https://doi.org/10.1016/j.biortech.2015.08.030

Singh, R., Kumar, M., Mittal, A., & Mehta, P. K. (2016). Microbial enzymes: Industrial progress in 21st century. *3 Biotech, 6*(2), 174. https://doi.org/10.1007/s13205-016-0485-8

Singh, A., Sharma, P., & Arora, A. (2018). Soybean meal as a substrate for microbial enzyme production: Advances and applications. *Bioresource Technology Reports, 4*, 50–56. https://doi.org/10.1016/j.biteb.2018.09.003

Singh, A., Sharma, P., & Gupta, R. (2018). Seasonal variability of agro-residues and its impact on microbial enzyme production. *Bioresource Technology, 269*, 450–457. https://doi.org/10.1016/j.biortech.2018.08.110

Singh, P., Kumar, V., & Sharma, A. (2025). Integrating microbial bioprocesses into circular bioeconomy frameworks for sustainable manufacturing. *Renewable and Sustainable Energy Reviews, 210*, 113841. https://doi.org/10.1016/j.rser.2024.113841

Soccol, C. R., da Costa, E. S. F., Letti, L. A. J., Karp, S. G., Woiciechowski, A. L., & de Souza Vandenberghe, L. P. (2017). Recent developments and innovations in solid-state fermentation. *Biotechnology Research and Innovation, 1*(1), 52–71. https://doi.org/10.1016/j.biori.2017.01.002

Sooch, B. S., Kaushik, S., & Singh, P. (2019). Waste valorization for enzyme production: Strategies and future perspectives. *Bioresource Technology Reports, 7*, 100244. https://doi.org/10.1016/j.biteb.2019.100244

Sooch, B. S., Kauldhar, B. S., & Randhawa, J. S. (2019). Cost-effective production of microbial enzymes from fruit and vegetable wastes. *Waste and Biomass Valorization, 10*(6), 1585–1595. https://doi.org/10.1007/s12649-018-0209-7

Soccol, C. R., Brar, S. K., & Pandey, A. (2021). *Biorefinery for sustainable production of fuels, chemicals, and proteins*. Springer Nature.

Sultana, N., Parveen, R., & Alam, M. (2024). Fruit and vegetable wastes as substrates for microbial enzyme synthesis: Recent advances. *Environmental Technology & Innovation, 35*, 103281. https://doi.org/10.1016/j.eti.2024.103281

Sun, Y., & Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: A review. *Bioresource Technology, 83*(1), 1–11. https://doi.org/10.1016/S0960-8524(01)00212-7

Wan, C., & Li, Y. (2012). Fungal pretreatment of lignocellulosic biomass. *Biotechnology Advances, 30*(6), 1447–1457. https://doi.org/10.1016/j.biotechadv.2012.03.003

Zhang, X., Xu, D., & Zhang, Y. (2019). Pretreatment strategies for improved microbial enzyme production from lignocellulosic biomass. *Renewable Energy, 138*, 691–701. https://doi.org/10.1016/j.renene.2019.01.103

Zhang, L., Wang, C., & Li, Y. (2024). Circular bioeconomy-driven enzyme bioprocessing: Integrating sustainability and innovation. *Journal of Cleaner Production, 438*, 140142. https://doi.org/10.1016/j.jclepro.2024.140142

Zhao, X., Zhang, L., & Liu, D. (2012). Biomass recalcitrance. Part I: The chemical compositions and physical structures affecting the enzymatic hydrolysis of lignocellulose. *Biofuels, Bioproducts and Biorefining, 6*(4), 465–482. https://doi.org/10.1002/bbb.1331

Zhao, Y., Liu, X., & Tang, J. (2024). Green solvent-based pretreatment of biomass for microbial enzyme production. *Industrial Crops and Products, 208*, 117523. https://doi.org/10.1016/j.indcrop.2024.117523